

A component of the mir-17-92 polycistronic oncomir promotes oncogene-dependent apoptosis.

Journal: Elife

Publication Year: 2013

Authors: Virginie Olive, Erich Sabio, Margaux J Bennett, Caitlin S De Jong, Anne Biton, James C McGann, Samantha K Greaney, Nicole M Sodik, Alicia Y Zhou, Asha Balakrishnan, Mona Foth, Micah A Luftig, Andrei Goga, Terence P Speed, Zhenyu Xuan, Gerard I Evan, Ying Wan, Alex C Minella, Lin He

PubMed link: 24137534

Funding Grants: The roles of non-coding RNAs in the self-renewal and differentiation of pluripotent stem cells

Public Summary:

mir-17-92, a potent polycistronic oncomir, encodes six mature miRNAs with complex modes of interactions. In the Emu-myc Burkitt's lymphoma model, mir-17-92 exhibits potent oncogenic activity by repressing c-Myc-induced apoptosis, primarily through its miR-19 components. Surprisingly, mir-17-92 also encodes the miR-92 component that negatively regulates its oncogenic cooperation with c-Myc. This miR-92 effect is, at least in part, mediated by its direct repression of Fbw7, which promotes the proteosomal degradation of c-Myc. Thus, overexpressing miR-92 leads to aberrant c-Myc increase, imposing a strong coupling between excessive proliferation and p53-dependent apoptosis. Interestingly, miR-92 antagonizes the oncogenic miR-19 miRNAs; and such functional interaction coordinates proliferation and apoptosis during c-Myc-induced oncogenesis. This miR-19:miR-92 antagonism is disrupted in B-lymphoma cells that favor a greater increase of miR-19 over miR-92. Altogether, we suggest a new paradigm whereby the unique gene structure of a polycistronic oncomir confers an intricate balance between oncogene and tumor suppressor crosstalk.

Scientific Abstract:

mir-17-92, a potent polycistronic oncomir, encodes six mature miRNAs with complex modes of interactions. In the Emu-myc Burkitt's lymphoma model, mir-17-92 exhibits potent oncogenic activity by repressing c-Myc-induced apoptosis, primarily through its miR-19 components. Surprisingly, mir-17-92 also encodes the miR-92 component that negatively regulates its oncogenic cooperation with c-Myc. This miR-92 effect is, at least in part, mediated by its direct repression of Fbw7, which promotes the proteosomal degradation of c-Myc. Thus, overexpressing miR-92 leads to aberrant c-Myc increase, imposing a strong coupling between excessive proliferation and p53-dependent apoptosis. Interestingly, miR-92 antagonizes the oncogenic miR-19 miRNAs; and such functional interaction coordinates proliferation and apoptosis during c-Myc-induced oncogenesis. This miR-19:miR-92 antagonism is disrupted in B-lymphoma cells that favor a greater increase of miR-19 over miR-92. Altogether, we suggest a new paradigm whereby the unique gene structure of a polycistronic oncomir confers an intricate balance between oncogene and tumor suppressor crosstalk.

DOI:<http://dx.doi.org/10.7554/eLife.00822.001>.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/component-mir-17-92-polycistronic-oncomir-promotes-oncogene-dependent>